

# Alcohol & Drug Dependence



## Follow-Up After Emergency Department Visit for Alcohol & Other Drug Abuse or Dependence (FUA)

Members 13 years of age and older who visited the emergency department (ED) with a principal diagnosis of alcohol or other drug (AOD) abuse or dependence and had a follow-up visit for AOD between January 1 and December 1 of the measurement year

Two rates are reported:

30-Day Follow-Up	Seven-Day Follow-Up
Members who had a follow-up visit with any practitioner within 30 days of the ED visit (31 total days)	Members who had a follow-up visit with any practitioner within seven days of the ED visit (eight total days)
<b>Note:</b> Can Include visits that occur on the date of the ED visit	<b>Note:</b> Can Include visits that occur on the date of the ED visit

Follow-Up Visits			
Visit Type	CPT®	HCPCS	POS
IET Stand Alone Visits with a principal diagnosis of AOD abuse or dependence	99201-99205, 99211-99215, 99241-99245, 99341-99343, 99345, 99347-99350, 99384-99385, 99394-99395, 99401-99402	H0001, H0015, H0034, H0035, H2000, H2011, H2014, H2015, H2016, H2017, H2019, H2020, T1015	
IET Visits Group 1 with a principal diagnosis of AOD abuse or dependence	90791, 90792, 90832, 90833, 90834, 90836, 90837, 90838, 90839, 90840, 90847, 90849, 90853		02, 03, 05, 07, 09, 11-20, 22, 33, 49, 50, 52, 53, 57, 58, 71, 72
IET Visits Group 2 with a principal diagnosis of AOD abuse or dependence	99221, 99222, 99223, 99231, 99232, 99233, 99238, 99251, 99252, 99253, 99254, 99255		02, 52, 53
Observation visit with a principal diagnosis of AOD abuse or dependence	99217, 99218, 99219, 99220		
A virtual check-in with a principal diagnosis of AOD abuse or dependence		G2012	

The educational material herein complies with accepted ICD-10 guidelines and is for general supplemental purposes only. The information herein is not guaranteed to be complete, free of errors, or the most current revision. It is the responsibility of the provider to document accurate and complete codes, clinical rationale, and medical services rendered to support appropriate ICD-10 code(s) according to official billing and coding guidelines, procedures, and regulations.

# Alcohol & Drug Dependence



## Coding & Documentation

The presence of a condition cannot be assumed even if evidence is present in the medical record. It requires the provider's clinical judgment and explicit diagnosis.

**Substance use disorder:** Patterned use of a substance in an amount or method causes significant impairment or distress.

**Physical Dependence:** A substance is needed to function normally; withdrawal symptoms would occur if the drug were stopped abruptly.

**Psychological Dependence:** Discontinued use of a substance results in emotional or behavioral disturbances.

## Diagnostic Criteria<sup>1</sup>

Patient must display two or more of the following symptoms in a 12-month period:

- Use of substance in larger doses or for a longer time period than intended
- Persistent desire or unsuccessful efforts to reduce or control use
- Abundance of time spent obtaining, using or recovering from the effects of a substance
- Strong cravings or desire to use
- Recurrent use resulting in failure to fulfill obligations at work, home or school
- Continued use even when persistent or recurrent social or interpersonal problems are caused by the effects
- Abandonment of important social, recreational or occupational activities because of use
- Recurrent use in physically hazardous situations
- Tolerance — the need for increased amount achieve desired effect; the diminished effect with continued use of same amount
- Withdrawal — characteristic withdrawal syndrome; use to relieve or avoid withdrawal

## ICD-10-CM Codes<sup>2</sup>

Mental and behavioral disorders due to psychoactive substance use, F10–F19

**Identify the substance** — each substance is addressed as a separate disorder:

- Alcohol
- Cocaine
- Inhalant
- Opioid
- Other stimulant (amphetamine, caffeine, etc.)
- Other (or unknown) psychoactive substance
- Cannabis
- Hallucinogen
- Sedative, hypnotic or anxiolytic
- Nicotine

1 American Psychiatric Association: Diagnostic and Statistical Manual of Mental Disorders: Diagnostic and Statistical Manual of Mental Disorders, Fifth Edition. Arlington, VA: American Psychiatric Association, 2013.

2 <https://www.cms.gov/medicare/icd-10/2021-icd-10-cm>

# Alcohol & Drug Dependence



## Coding Hierarchy

If Documented...	Then Code...
Use & abuse	Abuse
Use & dependence	Dependence
Abuse & dependence	Dependence
Use, abuse & dependence	Dependence

## Disorder Severity

Mild	2-3 symptoms
Moderate	4-5 symptoms
Severe	6+ symptoms

### Severe use disorder = addiction<sup>3</sup>

A substance addiction will likely require ongoing treatment for several years.

## “In Remission”

Disorder	Classification
Early remission	No signs/symptoms of disorder for 3-12 months
Sustained remission	No signs/symptoms of disorder for 12+ months
Abuse in remission	Mild use disorder in early or sustained remission
Dependence in remission	Moderate or severe use disorder in early or sustained remission

## Document to the highest known degree and code to the highest specificity.

Documentation should include, if applicable:

- Substance type
- Severity
- Association with intoxication or withdrawal
- Detailed clinical diagnosis
- Treatment plan/condition management
- Presence of induced manifestation or complication, such as:
  - Delirium
  - Perceptual disturbance
  - Anxiety
  - Sexual dysfunction
  - Sleep or mood disorder
  - Persisting amnesic disorder or dementia

<sup>3</sup> <https://www.addictionpolicy.org/post/dsm-5-facts-and-figures>

# Alcohol & Drug Dependence



## HEDIS® & Quality

### Initiation and Engagement of Alcohol & Other Drug Abuse or Dependence Treatment (IET)

Measure evaluates adolescent and adult members, ages 13 and older, with a new episode of alcohol or other drug (AOD) abuse or dependence who received the following between January 1 and November 14 of the measurement year

#### Two rates are reported:

Initiation of AOD Treatment	Engagement of AOD Treatment
Members who initiate treatment through an inpatient AOD admission, outpatient visit, intensive outpatient encounter, or partial hospitalization, telehealth, or medication treatment within 14 days of diagnosis	Members who initiated treatment and were engaged in ongoing AOD treatment within 34 days of the initiation visit. Engagement is at least one medication treatment event or two visits by AOD admission, outpatient visit, intensive outpatient encounter, or partial hospitalization, or telehealth.

#### Initiation of AOD Treatment

Submit claims with any diagnosis code to identify AOD with an IET treatment code as listed below.

**For all initiation events except medication treatment, initiation on the same day as the index episode start date (IESD) must be with different providers in order to count.**

Outpatient, intensive outpatient, partial hospitalization, observation, telehealth, or ED visit (not resulting in an inpatient stay)	The date of service
Inpatient stay or for detoxification that occurred during an inpatient stay	The date of discharge
Detoxification (other than detoxification that occurred during an inpatient stay)	The date of service
ED or observation visits that result in an inpatient stay	The date of the inpatient discharge
Direct transfers	The discharge date from the last admission

**For additional resources, contact our quality improvement team at [QI\\_AR\\_HEDIS@centene.com](mailto:QI_AR_HEDIS@centene.com)**

HEDIS Vol. 2 Technical Specifications by NCQA

**Note:** The information listed here is not all-inclusive and should be used only as reference. Please refer to the current ICD-10/CPT®/HCPCS coding and documentation guidelines at [cms.gov](http://cms.gov). HEDIS guidelines are available at [www.ncqa.org](http://www.ncqa.org)

# Alcohol & Drug Dependence



If the index episode was an inpatient discharge or ED/observation visit that resulted in an inpatient stay, then the inpatient stay is considered initiation of treatment and the member is compliant. If the index episode was *not* an inpatient discharge, then the member must initiate treatment on the IESD or in the 13 days following the IESD (14 total days). Any of the following combinations must meet criteria for initiation. ICD-10-CM codes for alcohol, opioid, or other drug abuse or dependence are listed on the next page.

Description	Coding		
An acute or non-acute inpatient admission <b>with</b> an alcohol, opioid or other drug abuse and/or dependence diagnosis on the discharge claim	<b>UBREV</b>		
	0100, 0101, 0110, 0111, 0112, 0113, 0114, 0116, 0117, 0118, 0119, 0120, 0121, 0122, 0123, 0124, 0126, 0127, 0128, 0129, 0130, 0131, 0132, 0133, 0134, 0136, 0137, 0138, 0139, 0140, 0141, 0142, 0143, 0144, 0146, 0147, 0148, 0149, 0150, 0151, 0152, 0153, 0154, 0156, 0157, 0158, 0159, 0160, 0164, 0167, 0169, 0170, 0171, 0172, 0173, 0174, 0179, 0190, 0191, 0192, 0193, 0194, 0199, 0200, 0201, 0202, 0203, 0204, 0206, 0207, 0208, 0209, 0210, 0211, 0212, 0213, 0214, 0219, 1000, 1001, 1002		
IET standalone visits <b>with</b> an alcohol, opioid or other drug abuse and/or dependence diagnosis	<b>CPT</b>	<b>HCPCS</b>	<b>UBREV</b>
	99201, 99202, 99203, 99204, 99205, 99211, 99212, 99213, 99214, 99215, 99241, 99242, 99243, 99244, 99245, 99341, 99342, 99343, 99345, 99347, 99348, 99349, 99350, 99384, 99385, 99386, 99387, 99394, 99395, 99396, 99397, 99401, 99402	H0001, H0015, H0034, H0035, H2000, H2011, H2014, H2015, H2016, H2017, H2019, H2020, H2035, T1015	0510, 0513, 0515, 0516, 0517, 0519, 0520, 0521, 0522, 0523, 0526, 0527, 0528, 0529, 0900, 0902, 0903, 0904, 0905, 0906, 0907, 0911, 0912, 0913, 0914, 0915, 0916, 0917, 0919, 0944, 0945, 0982, 0983
An observation <b>with</b> an alcohol, opioid or other drug abuse and/or dependence diagnosis	<b>CPT</b>		
	99217, 99218, 99219, 99220		
IET group 1 visits <b>with</b> POS and an alcohol, opioid or other drug abuse and/or dependence diagnosis	<b>CPT</b>	<b>Place of Service</b>	
	90791, 90792, 90832, 90833, 90834, 90836, 90837, 90838, 90839, 90840, 90847, 90849, 90853	02, 03, 05, 07, 09, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 22, 33, 49, 50, 52, 53, 57, 58, 71, 72	
IET group 2 visits <b>with</b> POS and an alcohol, opioid or other drug abuse and/or dependence diagnosis	<b>CPT</b>	<b>Place of Service</b>	
	99221, 99222, 99223, 99231, 99232, 99233, 99238, 99251, 99252, 99253, 99254, 99255	02, 52, 53	
A virtual check-in <b>with</b> an alcohol, opioid or other drug abuse and/ or dependence diagnosis	<b>HCPCS</b>		
	G2012		

# Alcohol & Drug Dependence



## Alcohol, Opioid, and Other Drug Abuse or Dependence ICD-10-CM Codes

F10.10, F10.120, F10.121, F10.129, F10.14, F10.150, F10.151, F10.159, F10.180, F10.181, F10.182, F10.188, F10.19, F10.20, F10.220, F10.221, F10.229, F10.230, F10.231, F10.232, F10.239, F10.24, F10.250, F10.251, F10.259, F10.26, F10.27, F10.280, F10.281, F10.282, F10.288, F10.29, F11.10, F11.120, F11.121, F11.122, F11.129, F11.14, F11.150, F11.151, F11.159, F11.181, F11.182, F11.188, F11.19, F11.20, F11.220, F11.221, F11.222, F11.229, F11.23, F11.24, F11.250, F11.251, F11.259, F11.281, F11.282, F11.288, F11.29, F12.10, F12.120, F12.121, F12.122, F12.129, F12.150, F12.151, F12.159, F12.180, F12.188, F12.19, F12.20, F12.220, F12.221, F12.222, F12.229, F12.23, F12.250, F12.251, F12.259, F12.280, F12.288, F12.29, F13.10, F13.120, F13.121, F13.129, F13.14, F13.150, F13.151, F13.159, F13.180, F13.181, F13.182, F13.188, F13.19, F13.20, F13.220, F13.221, F13.229, F13.230, F13.231, F13.232, F13.239, F13.24, F13.250, F13.251, F13.259, F13.26, F13.27, F13.280, F13.281, F13.282, F13.288, F13.29, F14.10, F14.120, F14.121, F14.122, F14.129, F14.14, F14.150, F14.151, F14.159, F14.180, F14.181, F14.182, F14.188, F14.19, F14.20, F14.220, F14.221, F14.222, F14.229, F14.23, F14.24, F14.250, F14.251, F14.259, F14.280, F14.281, F14.282, F14.288, F14.29, F15.10, F15.120, F15.121, F15.122, F15.129, F15.14, F15.150, F15.151, F15.159, F15.180, F15.181, F15.182, F15.188, F15.19, F15.20, F15.220, F15.221, F15.222, F15.229, F15.23, F15.24, F15.250, F15.251, F15.259, F15.280, F15.281, F15.282, F15.288, F15.29, F16.10, F16.120, F16.121, F16.122, F16.129, F16.14, F16.150, F16.151, F16.159, F16.180, F16.183, F16.188, F16.19, F16.20, F16.220, F16.221, F16.229, F16.24, F16.250, F16.251, F16.259, F16.280, F16.283, F16.288, F16.29, F18.10, F18.120, F18.121, F18.129, F18.14, F18.150, F18.151, F18.159, F18.17, F18.180, F18.188, F18.19, F18.20, F18.220, F18.221, F18.229, F18.24, F18.250, F18.251, F18.259, F18.27, F18.280, F18.288, F18.29, F19.10, F19.120, F19.121, F19.122, F19.129, F19.14, F19.150, F19.151, F19.159, F19.16, F19.17, F19.180, F19.181, F19.182, F19.188, F19.19, F19.20, F19.220, F19.221, F19.222, F19.229, F19.230, F19.231, F19.232, F19.239, F19.24, F19.250, F19.251, F19.259, F19.26, F19.27, F19.280, F19.281, F19.282, F19.288, F19.29

If the index episode was for a diagnosis of alcohol abuse or dependence, a medication treatment-dispensing event or medication treatment during a visit, then:

HCPSC Codes	ICD-10-CM Codes
J0570, J2315, Q9991, Q9992	F10.10, F10.120, F10.121, F10.129, F10.14, F10.150, F10.151, F10.159, F10.180, F10.181, F10.182, F10.188, F10.19, F10.20, F10.220, F10.221, F10.229, F10.230, F10.231, F10.232, F10.239, F10.24, F10.250, F10.251, F10.259, F10.26, F10.27, F10.280, F10.281, F10.282, F10.288, F10.29

Alcohol Use Disorder Treatment Medications	
Description	Prescription
Aldehyde dehydrogenase inhibitor	Disulfiram (oral)
Antagonist	Naltrexone (oral and injectable)
Other	Acamprosate (oral; delayed-release tablet)

If the index episode was for a diagnosis of opioid abuse or dependence, a medication treatment-dispensing event or medication treatment during a visit, then:

HCPSC Codes	ICD-10-CM Codes
J0570, J2315, Q9991, Q9992	F11.10, F11.120, F11.121, F11.122, F11.129, F11.14, F11.150, F11.151, F11.159, F11.181, F11.182, F11.188, F11.19, F11.20, F11.220, F11.221, F11.222, F11.229, F11.23, F11.24, F11.250, F11.251, F11.259, F11.281, F11.282, F11.288, F11.29

Opioid Use Disorder Treatment Medications	
Description	Prescription
Antagonist	Naltrexone (oral and injectable)
Partial agonist	<ul style="list-style-type: none"> <li>Buprenorphine (sublingual tablet, injection, implant)</li> <li>Buprenorphine/naloxone (sublingual tablet, buccal film, sublingual film)</li> </ul>

## Engagement of AOD Treatment

Submit claims with any diagnosis code to identify AOD with an IET treatment code as listed below:

- Two engagement visits can be on the same date of service, but they must be with different providers in order to count as two events. An engagement visit on the same date of service as an engagement medication treatment event meets criteria, and there is no requirement that they be with different providers.
- For members who initiated treatment via an inpatient admission, the 34-day period for engagement begins the day after discharge.
- If a member had an opioid treatment service that bills monthly, or if they had a visit that included medication administration beginning the day after the initiation encounter through 34 days following the initiation event, and if the IESD was a diagnosis of opioid abuse or dependence, then the member is compliant with engagement for AOD treatment.
- If a member's initiation of AOD treatment was a medication treatment event, and the member has two or more engagement events in which only one can be an engagement medication treatment event, beginning the day after the initiation encounter through 34 days following the initiation event (34 total days), then the member is compliant with engagement for AOD treatment.

**For additional resources, contact our quality improvement team at [QI\\_AR\\_HEDIS@centene.com](mailto:QI_AR_HEDIS@centene.com)**

HEDIS Vol. 2 Technical Specifications by NCQA

**Note:** The information listed here is not all-inclusive and should be used only as reference. Please refer to the current ICD-10/CPT®/HCPCS coding and documentation guidelines at [cms.gov](http://cms.gov). HEDIS guidelines are available at [www.ncqa.org](http://www.ncqa.org)

# Alcohol & Drug Dependence



Any of the following beginning the day after the initiation encounter through 34 days following the initiation event (34 total days) meet criteria for an engagement visit:

Description	Coding		
An acute or non-acute inpatient admission <b>with</b> an alcohol, opioid, or other drug abuse and/or dependence diagnosis on the discharge claim	<b>UBREV</b>		
	0100, 0101, 0110, 0111, 0112, 0113, 0114, 0116, 0117, 0118, 0119, 0120, 0121, 0122, 0123, 0124, 0126, 0127, 0128, 0129, 0130, 0131, 0132, 0133, 0134, 0136, 0137, 0138, 0139, 0140, 0141, 0142, 0143, 0144, 0146, 0147, 0148, 0149, 0150, 0151, 0152, 0153, 0154, 0156, 0157, 0158, 0159, 0160, 0164, 0167, 0169, 0170, 0171, 0172, 0173, 0174, 0179, 0190, 0191, 0192, 0193, 0194, 0199, 0200, 0201, 0202, 0203, 0204, 0206, 0207, 0208, 0209, 0210, 0211, 0212, 0213, 0214, 0219, 1000, 1001, 1002		
IET standalone visits <b>with</b> an alcohol, opioid, or other drug abuse and/or dependence diagnosis	<b>CPT</b>	<b>HCPCS</b>	<b>UBREV</b>
	99201, 99202, 99203, 99204, 99205, 99211, 99212, 99213, 99214, 99215, 99241, 99242, 99243, 99244, 99245, 99341, 99342, 99343, 99345, 99347, 99348, 99349, 99350, 99384, 99385, 99386, 99387, 99394, 99395, 99396, 99397, 99401, 99402	H0001, H0015, H0034, H0035, H2000, H2011, H2014, H2015, H2016, H2017, H2019, H2020, H2035, T1015	0510, 0513, 0515, 0516, 0517, 0519, 0520, 0521, 0522, 0523, 0526, 0527, 0528, 0529, 0900, 0902, 0903, 0904, 0905, 0906, 0907, 0911, 0912, 0913, 0914, 0915, 0916, 0917, 0919, 0944, 0945, 0982, 0983
An observation <b>with</b> an alcohol, opioid, or other drug abuse and/or dependence diagnosis	<b>CPT</b>		
	99217, 99218, 99219, 99220		
IET group 1 visits <b>with</b> POS <b>with</b> an alcohol, opioid, or other drug abuse and/or dependence diagnosis	<b>CPT</b>	<b>Place of Service</b>	
	90791, 90792, 90832, 90833, 90834, 90836, 90837, 90838, 90839, 90840, 90847, 90849, 90853	02, 03, 05, 07, 09, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 22, 33, 49, 50, 52, 53, 57, 58, 71, 72	
IET group 2 visits <b>with</b> POS <b>with</b> an alcohol, opioid, or other drug abuse and/or dependence diagnosis	<b>CPT</b>	<b>Place of Service</b>	
	99221, 99222, 99223, 99231, 99232, 99233, 99238, 99251, 99252, 99253, 99254, 99255	02, 52, 53	
A virtual check-in <b>with</b> an alcohol, opioid, or other drug abuse and/or dependence diagnosis	<b>HCPCS</b>		
	G2012		



# Alcohol & Drug Dependence



## Alcohol, Opioid, and Other Drug Abuse and Dependence ICD-10-CM Codes

F10.10, F10.120, F10.121, F10.129, F10.14, F10.150, F10.151, F10.159, F10.180, F10.181, F10.182, F10.188, F10.19, F10.20, F10.220, F10.221, F10.229, F10.230, F10.231, F10.232, F10.239, F10.24, F10.250, F10.251, F10.259, F10.26, F10.27, F10.280, F10.281, F10.282, F10.288, F10.29, F11.10, F11.120, F11.121, F11.122, F11.129, F11.14, F11.150, F11.151, F11.159, F11.181, F11.182, F11.188, F11.19, F11.20, F11.220, F11.221, F11.222, F11.229, F11.23, F11.24, F11.250, F11.251, F11.259, F11.281, F11.282, F11.288, F11.29, F12.10, F12.120, F12.121, F12.122, F12.129, F12.150, F12.151, F12.159, F12.180, F12.188, F12.19, F12.20, F12.220, F12.221, F12.222, F12.229, F12.23, F12.250, F12.251, F12.259, F12.280, F12.288, F12.29, F13.10, F13.120, F13.121, F13.129, F13.14, F13.150, F13.151, F13.159, F13.180, F13.181, F13.182, F13.188, F13.19, F13.20, F13.220, F13.221, F13.229, F13.230, F13.231, F13.232, F13.239, F13.24, F13.250, F13.251, F13.259, F13.26, F13.27, F13.280, F13.281, F13.282, F13.288, F13.29, F14.10, F14.120, F14.121, F14.122, F14.129, F14.14, F14.150, F14.151, F14.159, F14.180, F14.181, F14.182, F14.188, F14.19, F14.20, F14.220, F14.221, F14.222, F14.229, F14.23, F14.24, F14.250, F14.251, F14.259, F14.280, F14.281, F14.282, F14.288, F14.29, F15.10, F15.120, F15.121, F15.122, F15.129, F15.14, F15.150, F15.151, F15.159, F15.180, F15.181, F15.182, F15.188, F15.19, F15.20, F15.220, F15.221, F15.222, F15.229, F15.23, F15.24, F15.250, F15.251, F15.259, F15.280, F15.281, F15.282, F15.288, F15.29, F16.10, F16.120, F16.121, F16.122, F16.129, F16.14, F16.150, F16.151, F16.159, F16.180, F16.183, F16.188, F16.19, F16.20, F16.220, F16.221, F16.229, F16.24, F16.250, F16.251, F16.259, F16.280, F16.283, F16.288, F16.29, F18.10, F18.120, F18.121, F18.129, F18.14, F18.150, F18.151, F18.159, F18.17, F18.180, F18.188, F18.19, F18.20, F18.220, F18.221, F18.229, F18.24, F18.250, F18.251, F18.259, F18.27, F18.280, F18.288, F18.29, F19.10, F19.120, F19.121, F19.122, F19.129, F19.14, F19.150, F19.151, F19.159, F19.16, F19.17, F19.180, F19.181, F19.182, F19.188, F19.19, F19.20, F19.220, F19.221, F19.222, F19.229, F19.230, F19.231, F19.232, F19.239, F19.24, F19.250, F19.251, F19.259, F19.26, F19.27, F19.280, F19.281, F19.282, F19.288, F19.29

If the index episode was for a diagnosis of opioid abuse or dependence during an opioid treatment service, then:

## ICD-10-CM Codes

F11.10, F11.120, F11.121, F11.122, F11.129, F11.14, F11.150, F11.151, F11.159, F11.181, F11.182, F11.188, F11.19, F11.20, F11.220, F11.221, F11.222, F11.229, F11.23, F11.24, F11.250, F11.251, F11.259, F11.281, F11.282, F11.288, F11.29

**For additional resources, contact our quality improvement team at [QI\\_AR\\_HEDIS@centene.com](mailto:QI_AR_HEDIS@centene.com)**

HEDIS Vol. 2 Technical Specifications by NCQA

**Note:** The information listed here is not all-inclusive and should be used only as reference. Please refer to the current ICD-10/CPT®/HCPCS coding and documentation guidelines at [cms.gov](https://www.cms.gov). HEDIS guidelines are available at [www.ncqa.org](https://www.ncqa.org)

# Alcohol & Drug Dependence



If the IESD diagnosis was a diagnosis of alcohol abuse or dependence, one or more medication treatment-dispensing events, or medication treatment during a visit, then beginning the day after the initiation encounter through 34 days following the initiation event (34 total days), this meets criteria for alcohol abuse and dependence treatment.

HCPCS Codes	ICD-10-CM Codes
J0570, J2315, Q9991, Q9992	F10.10, F10.120, F10.121, F10.129, F10.14, F10.150, F10.151, F10.159, F10.180, F10.181, F10.182, F10.188, F10.19, F10.20, F10.220, F10.221, F10.229, F10.230, F10.231, F10.232, F10.239, F10.24, F10.250, F10.251, F10.259, F10.26, F10.27, F10.280, F10.281, F10.282, F10.288, F10.29

Alcohol Use Disorder Treatment Medications	
Description	Prescription
Aldehyde dehydrogenase inhibitor	Disulfiram (oral)
Antagonist	Naltrexone (oral and injectable)
Other	Acamprosate (oral; delayed-release tablet)

If the IESD was a diagnosis of opioid abuse or dependence, one or more medication-dispensing events, or medication treatment during a visit, then beginning the day after the initiation encounter through 34 days following the initiation event (34 total days), this meets criteria for opioid abuse and dependence treatment.

HCPCS Codes	ICD-10-CM Codes
J0570, J2315, Q9991, Q9992	F11.10, F11.120, F11.121, F11.122, F11.129, F11.14, F11.150, F11.151, F11.159, F11.181, F11.182, F11.188, F11.19, F11.20, F11.220, F11.221, F11.222, F11.229, F11.23, F11.24, F11.250, F11.251, F11.259, F11.281, F11.282, F11.288, F11.29

Opioid Use Disorder Treatment Medications	
Description	Prescription
Antagonist	Naltrexone (oral and injectable)
Partial agonist	<ul style="list-style-type: none"> <li>Buprenorphine (sublingual tablet, injection, implant)</li> <li>Buprenorphine/naloxone (sublingual tablet, buccal film, sublingual film)</li> </ul>

**For additional resources, contact our quality improvement team at [QI\\_AR\\_HEDIS@centene.com](mailto:QI_AR_HEDIS@centene.com)**

HEDIS Vol. 2 Technical Specifications by NCQA

**Note:** The information listed here is not all-inclusive and should be used only as reference. Please refer to the current ICD-10/CPT®/HCPCS coding and documentation guidelines at cms.gov. HEDIS guidelines are available at [www.ncqa.org](http://www.ncqa.org)

# Alcohol & Drug Dependence



## Use of Opioid at High Dosage (HDO)

Members 18 years and older who received prescription opioids at a high dosage (average morphine milligram equivalent dose (MME)  $\geq$  90) for 15 days or more during the measurement year

**Note:** A lower rate indicates better performance.

Opioid Medications					
Type of Opioid	Medication Lists & Strengths	MME Conversion Factor <sup>4</sup>	Type of Opioid	Medication Lists & Strengths	MME Conversion Factor <sup>4</sup>
Butorphanol	Butorphanol 10 MGPML	7	Fentanyl transdermal film/patch (mcg/hr)	Fentanyl 12 MCGPH, 12 MCGPH, 25 MCGPH, 37.5 MCGPH, 50 MCGPH, 62.5 MCGPH, 75 MCGPH, 87.5 MCGPH, 100 MCGPH	7.2
Codeine	Codeine Sulfate 15mg, 30mg, 60mg	0.15	Hydrocodone	Hydrocodone 10mg, 15mg, 20mg, 30mg, 40mg, 60mg, 80mg, 100mg, 120mg	1
Codeine	Acetaminophen-Codeine 2.4 MGPML, 15 mg, 30 mg, 60 mg	0.15	Hydrocodone	Acetaminophen-Hydrocodone .5 MGPML, .67 MGPML, 2.5mg, 5mg, 7.5mg, 7.5 MGPML, 10mg	1
Codeine	Acetaminophen-Butalbital-Caffeine-Codeine 30 mg	0.15	Hydrocodone	Hydrocodone-Ibuprofen 2.5mg, 5mg, 7.5mg, 10 mg	1
Codeine	Aspirin-Butalbital-Caffeine-Codeine 30 mg	0.15	Hydromorphone	Hydromorphone 1 MGPML, 2 mg, 3 mg, 4 mg, 8 mg, 12 mg, 16 mg, 32 mg	4
Codeine	Aspirin-Carisoprodol-Codeine 16 mg	0.15	Meperidine	Meperidine 10 MGPML, 50 mg, 100 mg	0.1
Dihydrocodeine	Acetaminophen-Caffeine-Dihydrocodeine 16 mg	0.15	Methadone	Methadone 1 MGPML, 2 MGPML, 5 mg, 10 mg, 10 MGPML, 40 mg	3
Dihydrocodeine	Aspirin-Caffeine-Dihydrocodeine 16 mg	0.25	Morphine	Morphine .4 MGPML, 2 MGPML, 4 MGPML, 5 mg, 10 mg, 15 mg, 20 MGPML, 20 mg, 30 mg, 40 mg, 45 mg, 50 mg, 60 mg, 70 mg, 75 mg, 80 mg, 90 mg, 100 mg, 120 mg, 130 mg, 150 mg, 200 mg	1
Fentanyl buccal or sublingual tablet, transmucosal lozenge (mcg)	Fentanyl 100 mcg, 200 mcg, 300 mcg, 400 mcg, 600 mcg, 800 mcg, 1200 mcg, 1600 mcg	0.13	Morphine	Opium Tincture .4 MGPML, 10 MGPML	1

<sup>4</sup> Determine the average MME. Sum the total daily MME for the treatment period and divide by the number of days in the treatment period. Members whose average MME was  $\geq$ 90 meet the numerator criteria.

# Alcohol & Drug Dependence



Opioid Medications					
Type of Opioid	Medication Lists & Strengths	MME Conversion Factor <sup>4</sup>	Type of Opioid	Medication Lists & Strengths	MME Conversion Factor <sup>4</sup>
Fentanyl oral spray (mcg)	Oral Fentanyl 100 MCGPS, 200 MCGPS, 400 MCGPS, 600 MCGPS, 800 MCGPS	0.18	Opium	Belladonna Opium 30 mg, 60 mg	1.5
Fentanyl nasal spray (mcg)	Nasal Fentanyl 100 MCGPS, 300 MCGPS, 400 MCGPS	0.16	Oxycodone	Oxycodone 1 MGPML, 5 mg Oxycodone 7.5 mg, 9 mg, 10 mg, 13.5 mg, 15 mg, 18 mg, 20 mg, 20 MGPML, 27 mg, 30 mg, 36 mg, 40 mg, 60 mg, 80 mg	0.37
Oxycodone	Acetaminophen-Oxycodone 2.5 mg, 5 mg, 7.5 mg, 10 mg	1.5	Pentazocine	Naloxone-Pentazocine 50 mg	0.4
Oxycodone	Aspirin-Oxycodone 4.8355 mg	1.5	Tapentadol	Tapentadol 50 mg, 75 mg, 100 mg, 150 mg, 200 mg, 250 mg	0.1
Oxycodone	Ibuprofen-Oxycodone 5 mg	1.5	Tramadol	Tramadol 50 mg, 100 mg, 150 mg, 200 mg, 300 mg	0.1
Oxymorphone	Oxymorphone 5 mg, 7.5 mg, 10 mg, 15 mg, 20 mg, 30 mg, 40 mg	3	Tramadol	Acetaminophen-Tramadol 37.5 mg	

# Alcohol & Drug Dependence



## Pharmacotherapy for Opioid Use Disorder (POD)

New opioid use disorder (OUD) pharmacotherapy events with OUD pharmacotherapy for 180 or more days among members ages 16 and older with a diagnosis of OUD between July 1 of the year prior to the measurement year and June 30 of the measurement year

**Note:** 180 or more days without a gap in treatment of 8 or more consecutive days

Opioid Use Disorder Treatment Medications			
Description	Prescription	HCPCS Code(s)	Supply
Antagonist	Naltrexone (oral)	N/A	N/A
Antagonist	Naltrexone (injection)	J2315	31-day supply
Partial agonist	Buprenorphine (sublingual tablet)	J0570	One-day supply
Partial agonist	Buprenorphine (injection)	Q9991, Q9992	31-day supply
Partial agonist	Buprenorphine (implant)	J0570	180-day supply

**Note:** Methadone is not included on the medication lists for this measure. Methadone for OUD administered or dispensed by federally certified opioid treatment programs (OTP) is billed on a medical claim. A pharmacy claim for methadone would be indicative of treatment for pain rather than OUD.

# Alcohol & Drug Dependence



## Use of Opioids From Multiple Providers (UOP)

Members 18 years and older who are receiving prescription opioids for 15 days or more during the measurement year, and who received opioids from multiple providers

**Note:** A lower rate indicates better performance for all three rates

Three rates are reported:

Multiple Prescribers	Multiple Prescribers and Pharmacies	Multiple Pharmacies
Members receiving prescriptions for opioids from four or more different prescribers during the measurement year	Members receiving prescriptions for opioids from four or more different prescribers <b>and</b> four or more different pharmacies during the measurement year (the proportion of members who are numerator-compliant for both the Multiple Prescribers and Multiple Pharmacies rates)	Members receiving prescriptions for opioids from four or more different pharmacies during the measurement year

Opioid Medications			
Prescriptions		Medication Lists	
Buprenorphine (transdermal patch and buccal film)	Buprenorphine	Methadone	Methadone
Butorphanol	Butorphanol	Morphine	Morphine
Codeine	Acetaminophen-Butalbital-Caffeine-Codeine, Acetaminophen-Codeine, Aspirin-Butalbital-Caffeine-Codeine, Aspirin-Carisoprodol-Codeine, Codeine Sulfate	Opium	Belladonna-Opium, Opium
Dihydrocodeine	Acetaminophen-Caffeine-Dihydrocodeine, Aspirin-Caffeine-Dihydrocodeine	Oxycodone	Acetaminophen-Oxycodone, Aspirin-Oxycodone, Ibuprofen-Oxycodone, Oxycodone
Fentanyl	Fentanyl	Oxymorphone	Oxymorphone
Hydrocodone	Acetaminophen-Hydrocodone, Hydrocodone, Hydrocodone-Ibuprofen	Pentazocine	Naloxone-Pentazocine
Hydromorphone	Hydromorphone	Tapentadol	Tapentadol
Levorphanol	Levorphanol	Tramadol	Acetaminophen-Tramadol, Tramadol
Meperidine	Meperidine		
	Benzhydrocodone	Acetaminophen Benzhydrocodone	